

Stronger steerability criterion for more uncertain continuous variable systems

Priyanka Chowdhury,^{1,*} Tanumoy Pramanik,^{2,†} and A. S. Majumdar^{1,‡}

¹*S. N. Bose National Centre for Basic Sciences,*

Block JD, Sector III, Salt Lake, Kolkata 700098, India

²*LTCI, Télécom ParisTech, 23 avenue d'Italie, 75214 Paris CEDEX 13, France*

(Dated: October 19, 2015)

We derive a fine-grained uncertainty relation for the measurement of two incompatible observables on a single quantum system of continuous variables, and show that continuous variable systems are more uncertain than discrete variable systems. Using the derived fine-grained uncertainty relation, we formulate stronger steering criterion that is able to reveal the steerability of N00N states that has hitherto not been possible using other criteria. We further obtain a monogamy relation for our steering inequality which leads to an, in principle, improved lower bound on the secret key rate of a one-sided device independent quantum key distribution protocol for continuous variables.

PACS numbers: 03.65.Ud, 42.50.Xa

I. INTRODUCTION

First, Heisenberg in his seminal paper [1] noted that two non-commuting observables in quantum mechanics could not be measured with arbitrary precision. The uncertainty principle introduces a sharp distinction between quantum and classical physics. Subsequently, a number of improved uncertainty relations have been provided [2–6]. The presence of uncertainty relations endows quantum mechanics with significant advantages over classical mechanics for performing different information processing tasks. Various versions of uncertainty relations have been used to detect entanglement [5, 7], to classify mixedness of states [8], to categorize different physical theories according to their strength of nonlocality [6, 9], and to bound information leakage in key distribution [10–15].

Uncertainty relations are linked directly to the ability of quantum states to enable steering. The phenomenon of quantum steering [16] emerges from the EPR paradox [17] that was first formulated for experimental realization [18] based on the Heisenberg uncertainty relation. Later, steering criteria [19] have been formulated using entropic uncertainty relations [3, 4], which reveal steering in states possessing higher order correlations [20]. Experimental demonstrations of steering have been performed beginning with the original demonstration of the EPR paradox [21], as well as using different settings [22] and loophole free arrangements [23]. Nonetheless, there exist states such as the N00N state which fail to display steering using the existing steering criteria for higher values of N in spite of violating Bell-type inequalities [24]. Such a feature calls for further improved steerability conditions since steering lies between entanglement and nonlocality in the hierarchy [25] of quantum correlations.

The tightest steering inequality in discrete variable sys-

tems is obtained [14] through the application of the fine-grained uncertainty relation (FUR) [6]. Fine-graining makes it possible to distinguish the uncertainty inherent in obtaining any particular combination of outcomes for different measurements. Application of fine-graining leads to discrimination between various physical theories based on the strength of nonlocality [6, 9]. It also provides an optimal lower bound of entropic uncertainty in presence of quantum memory [13]. In the present work we derive a FUR for continuous variables. Using the derived FUR we formulate a continuous variable fine-grained steering criterion, and show that this criterion is monogamous, i.e., Alice's steerability of Bob to a particular state chosen randomly from the eigenstates of two incompatible observables restricts the control of Bob's system by any eavesdropper. The security of one-sided device independent quantum key distribution (1s-DIQKD) [12] protocols rely on demonstration of steering. Here we show the possibility of obtaining of an improved lower bound on the key rate of a 1s-DIQKD protocol using our stronger steering inequality for continuous variables.

A comparison of the steering inequality derived here with the fine-grained steering inequality for discrete variables [14] is facilitated by using the Wigner function for computing various probabilities associated with continuous variables. There exists an analogy between the measurement of spin-1/2 projectors and the parity operator, since the measurement outcomes for both are dichotomic. It is well known [26] that the Wigner function expressed as an expectation value of a product of displaced parity operators can be used to derive Bell-CHSH inequalities [24] for continuous variables. Here we use the Wigner function formalism [26] to derive a steering inequality for continuous variables. We show that fine-graining leads to a novel manifestation of higher uncertainty in continuous variable systems, thereby enhancing, in principle, the key rate of a 1s-DIQKD protocol.

*Electronic address: priyanka@bose.res.in

†Electronic address: Pramanik@telecom-paristech.fr

‡Electronic address: archan@bose.res.in

II. FINE-GRAINED UNCERTAINTY RELATION FOR CONTINUOUS VARIABLES

Let us begin with a brief description of EPR-steering by considering the following game [25]. Alice prepares two systems A and B in the state ρ_{AB} and sends the system B to Bob. Alice's task is to convince Bob that the prepared state ρ_{AB} is entangled. Bob does not trust Alice, but he trusts that he receives a quantum system B . He is convinced only when the correlation between his outcome b for the measurement chosen randomly from the set $\mathcal{B} \in \{\beta_1, \beta_2\}$ and Alice's outcome a for the measurement chosen randomly from the set $\mathcal{A} \in \{\alpha_1, \alpha_2\}$ can not be explained by a local hidden state (LHS) model, *i.e.*,

$$P(a_{\mathcal{A}}, b_{\mathcal{B}}) = \sum_{\lambda} P(\lambda) P(a_{\mathcal{A}}|\lambda) P_Q(b_{\mathcal{B}}|\lambda), \quad (1)$$

where $P(\lambda)$ is a positive valued distribution over a set of hidden variables λ , and P_Q denotes probability of an outcome obtained from a quantum measurement. We assume here that Alice knows about Bob's set of observables.

Now, from Eq. (1) it is easy to derive the relation

$$P(b_{\mathcal{B}}|a_{\mathcal{A}}) \leq \max_{\lambda} [P_Q(b_{\mathcal{B}}|\lambda)] = P_Q(b_{\mathcal{B}}|\lambda_{\max}) \quad (2)$$

using $\sum_i x_i y_i \leq \max_i [x_i] \sum_i y_i \forall \{x_i, y_i\} \geq 0$. Next, using $\sum_i x_i y_i \geq \min_i [x_i] \sum_i y_i \forall \{x_i, y_i\} \geq 0$, we get

$$P(b_{\mathcal{B}}|a_{\mathcal{A}}) \geq \min_{\lambda} [P_Q(b_{\mathcal{B}}|\lambda)] = P_Q(b_{\mathcal{B}}|\lambda_{\min}). \quad (3)$$

Combining the relations (2) and (3), the sum of conditional probability distributions according to the LHS model is bounded by

$$\begin{aligned} & \min_{\beta_1, \beta_2} [P_Q(b_{\beta_1}|\lambda_{\min}) + P_Q(b_{\beta_2}|\lambda_{\min})] \\ & \leq P(b_{\beta_1}|a_{\alpha_1}) + P(b_{\beta_2}|a_{\alpha_2}) \leq \\ & \max_{\beta_1, \beta_2} [P_Q(b_{\beta_1}|\lambda_{\max}) + P_Q(b_{\beta_2}|\lambda_{\max})]. \end{aligned} \quad (4)$$

We formulate the FUR for continuous variables in order to obtain the bounds of the algebraic inequality (4).

The concept of fine-graining in uncertainty relations was first introduced by Oppenheim and Wehner [6] to explain the failure of quantum theory to exhibit the full non-local strength allowed by no-signaling theory. They bound an event (which is defined by the outcomes chosen using imposed restrictions or conditions) by its minimum possible uncertainty, or maximum possible certainty, for two incompatible observables. For qubit systems a game [6] is considered in which a binary question $q \in \{0, 1\}$ is given randomly to a player Bob who measures σ_z (σ_x) when $q = 0$ (1) is received. Here, the average uncertainty of getting spin up outcome (labeled by " $b = 0$ ") irrespective of the given question (or indeed, the average certainty) where the average is taken over all possible choice of measurements, is bounded by

$$\frac{1}{2} - \frac{1}{2\sqrt{2}} \leq \frac{1}{2} [P(b_{\sigma_z} = 0) + P(b_{\sigma_x} = 0)] \leq \frac{1}{2} + \frac{1}{2\sqrt{2}}, \quad (5)$$

where the equalities occur for maximally certain states. Here they are the eigenstates of the observables $(\sigma_z + \sigma_x)/\sqrt{2}$ and $(\sigma_z - \sigma_x)/\sqrt{2}$ for the upper and lower bounds, respectively. The bounds remain the same for the spin down outcome ($b = 1$). Using the uncertainty relation (5), the inequality (4) is bounded by $[1 - \frac{1}{\sqrt{2}}, 1 + \frac{1}{\sqrt{2}}]$. Hence, in discrete variable systems, the shared state is steerable if the value of $\frac{1}{2} [P(b_{\sigma_z} = 0) + P(b_{\sigma_x} = 0)]$ lies outside the above range [14], where Alice has prior knowledge of Bob's measurement settings.

In continuous variable systems, Bell's inequality is shown to be violated using the Wigner function formalism [26]. Fine-graining connects uncertainty with nonlocality, and hence for a given Bell-CHSH inequality one can formulate a FUR for a bipartite system [6]. Similar considerations hold true for single particle quantum systems, thus making it possible to construct a FUR for single systems using the Wigner distribution representing the average of displaced parity measurement. Let us here label the outcome of even parity measurement by "0". The corresponding projection operator is given by

$$\Pi^+(\beta) = \mathcal{D}(\beta) \left(\sum_{n=0}^{\infty} |2n\rangle \langle 2n| \right) \mathcal{D}^\dagger(\beta), \quad (6)$$

where $\mathcal{D}(\beta) (= \exp(\beta \hat{b}^\dagger - \beta^* \hat{b}))$, is the displacement operator with coherent displacement β , and $\hat{b}$ and $\hat{b}^\dagger$ the annihilation and creation operators, respectively. Similarly, the projection operator corresponding to the odd parity measurement outcome labeled by "1" is given by

$$\Pi^-(\beta) = \mathcal{D}(\beta) \left(\sum_{n=0}^{\infty} |2n+1\rangle \langle 2n+1| \right) \mathcal{D}^\dagger(\beta). \quad (7)$$

The observable associated with the Wigner function is given by $\hat{\mathcal{W}}(\beta) = \Pi^+(\beta) - \Pi^-(\beta)$ which can be realized using detectors with the capability of distinguishing the number of absorbed photons [26]. We will take β 's to be real displacements in the rest of this work.

The average certainty of the parity measurement outcome b over displacements α and β is given by $\frac{1}{2} [P(b_\alpha) + P(b_\beta)]$. As in the case of discrete variables, the average certainty here too is bounded by the minimum uncertainty states. In continuous variable systems it is well known that the coherent states

$$|\gamma\rangle = \exp\left[-\frac{|\gamma|^2}{2}\right] \sum_{m=0}^{\infty} \frac{\gamma^m}{\sqrt{m!}} |m\rangle \quad (8)$$

correspond to the minimum uncertainty states in phase-space. Therefore, we obtain the fine-grained bounds on $\frac{1}{2} [P(b_\alpha) + P(b_\beta)]$ using the coherent states.

For even parity measurement ($b = 0$) at the displace-

ment chosen from $\{\alpha, \beta\}$, the average certainty becomes

$$\begin{aligned} \frac{1}{2}[P(b_\alpha = 0) + P(b_\beta = 0)] &= \langle \gamma | \frac{\Pi^+(\alpha) + \Pi^+(\beta)}{2} | \gamma \rangle \\ &= \frac{1}{2} (\exp[-|\gamma - \alpha|^2] \cosh[|\gamma - \alpha|^2] \\ &\quad + \exp[-|\gamma - \beta|^2] \cosh[|\gamma - \beta|^2]), \end{aligned} \quad (9)$$

where, similar to the displacements β and α , we choose γ also to be real. Note here that the condition $\alpha = \beta \rightarrow \gamma$ needs to be excluded in order to ensure that the average certainty of getting even parity for the zero photon state does not always stay close to 1, (this is similar to getting, say, spin up outcome in discrete variables for spin measurements along directions $\hat{i}$ and $\hat{j}$ when $\hat{i} \rightarrow \hat{j}$). For simplification we henceforth set $\alpha = -\beta$, and compute the probability distribution $[P(b_\beta = 0) + P(b_{-\beta} = 0)]$. The validity of the FUR (4) is ensured by avoiding the region where both $\beta \rightarrow 0$ and $\gamma \rightarrow 0$, as shown in the Fig. (1). The probability distribution $[P(b_\beta = 0) + P(b_{-\beta} = 0)]$ is bounded by $[\frac{1}{2}, \frac{3}{4}]$, where the maximum occurs for $\beta = \gamma$. Similarly, the average certainty of odd parity measurements for the displacements β and $-\beta$ is bounded by

$$\begin{aligned} \frac{1}{2}[P(b_\beta = 1) + P(b_{-\beta} = 1)] &= \langle \gamma | \frac{\Pi^-(\beta) + \Pi^-(-\beta)}{2} | \gamma \rangle \\ &= \frac{1}{2} (\exp[-(\gamma - \beta)^2] \sinh[(\gamma - \beta)^2] \\ &\quad + \exp[-(\gamma + \beta)^2] \sinh[(\gamma + \beta)^2]). \end{aligned} \quad (10)$$

Hence, one gets $\frac{1}{4} \leq \frac{1}{2}[P(b_\beta = 1) + P(b_{-\beta} = 1)] \leq \frac{1}{2}$, except at $\gamma \rightarrow 0$ & $\beta \rightarrow 0$ for which the probability of getting odd counts for the zero photon state approaches zero (See Fig. (1)).

Combining the cases of odd and even parities, one finds that the FUR bounds the certainty for the measurement of two incompatible continuous variables observables by

$$\frac{1}{4} \leq \frac{1}{2}[P(b_\beta) + P(b_{-\beta})] \leq \frac{3}{4}. \quad (11)$$

In the Fig. (1), we plot the infimum value of $\frac{1}{2}[P(b_\beta) + P(b_{-\beta})]$ with γ . When $\beta \rightarrow 0$ the FUR (11) remains valid for $|\gamma|^2 \geq 1$, i.e., for a source with at least a single average photon number (we only need to avoid the situation when $\gamma \rightarrow 0$ & $\beta \rightarrow 0$ simultaneously). Note that the range of certainty in discrete variable systems given by Eq.(5) [14] is higher than that in continuous variable systems. This feature reflecting more uncertainty helps to improve the secret key rate, since higher uncertainty enables less information to the eavesdropper.

Now, using the FUR (11), the steering inequality (4) becomes

$$\frac{1}{4} \leq \frac{1}{2}[P(b_\beta|a_{\alpha_1}) + P(b_{-\beta}|a_{\alpha_2})] \leq \frac{3}{4}. \quad (12)$$

The violation of the inequality (12) indicates that the measurement correlations are unable to be explained with the help of a LHS model, i.e., the given state is steerable. Next, we provide an application of our derived steering inequality for N00N states.




FIG. 1: Plot of $\frac{1}{2} \inf_\beta [P(b_\beta = 0) + P(b_{-\beta} = 0)]$ with γ . The solid curve is for $\frac{1}{2} \max_\beta [P(b_\beta = 0) + P(b_{-\beta} = 0)]$, and the dashed curve is for $\frac{1}{2} \min_\beta [P(b_\beta = 1) + P(b_{-\beta} = 1)]$. The solid and dashed lines correspond, respectively, to the upper and lower bounds of certainty in regions of validity of the FUR (11).

III. STEERABILITY OF NOON STATES

N00N states [27] are regarded to be of high utility in quantum metrology for making precise interferometric measurements. Such a state is a maximally path-entangled two-mode number state of continuous variables, given by [28]

$$|N00N\rangle = \frac{1}{\sqrt{2}}(|N, 0\rangle - |0, N\rangle), \quad (13)$$

where N is the number of photons either in the first or the second mode. These states have been experimentally realized up to $N = 5$ [29]. The entanglement of N00N states given in terms of their logarithmic negativity is independent of the value of N , and Bell's inequality is maximally violated for all N [30]. However, they do not violate the entropic steering inequality for $N \geq 2$ [20]. We show now that such states are steerable for $N \geq 2$ using our derived steering inequality.

Considering the cases for N even and odd separately, we find that for the former the maximum violation of the inequality (12) on the upper side occurs when $\frac{1}{2}[P(b_\beta|a_{\alpha_1}) + P(b_{-\beta}|a_{\alpha_2})] = 1$ for the choices of the parameters given by $\{b = 0, a = 0\}$, $\{b = 0, a = 1\}$. Similarly, the maximum violation on the lower side occurs when $\frac{1}{2}[P(b_\beta|a_{\alpha_1}) + P(b_{-\beta}|a_{\alpha_2})] = 0$ for the choices of the parameters give by $\{b = 1, a = 0\}$, $\{b = 1, a = 1\}$. When N is odd, the maximum violations on the upper side are 1 for the choices $\{b = 0, a = 1\}$, $\{b = 1, a = 0\}$, and on the lower side are 0 for the choices $\{b = 0, a = 0\}$, $\{b = 1, a = 1\}$. In the Fig. (2), we plot of the quantity $\frac{1}{2}[P(b_\beta = 0|a_\alpha = 1) + P(b_{-\beta} = 0|a_{-\alpha} = 1)]$ versus β and α for $N = 2, 4$, and 6. One sees that the violation of the steering inequality occurs maximally for $N \geq 2$ in the region $|\beta| \rightarrow 0$. The condition for validity of our FUR, *viz.*, $|\gamma| \geq 1$ when $\beta \rightarrow 0$, is ensured since the average photon number ($N/2$ here) is greater than 1.




FIG. 2: Coloronline. The variation of $\frac{1}{2}[P(b_\beta = 0|a_\alpha = 1) + P(b_{-\beta} = 0|a_{-\alpha} = 1)]$ with respect to β and α for three different values of N . (i.) The red colored curve corresponds to the value $N = 2$; (iii) the green colored curve is for $N = 4$; and the blue colored curve is for $N = 6$.

IV. SECURITY OF KEY GENERATION

Steering finds direct applicability in demonstration of security of quantum key distribution (QKD). The goal of any QKD protocol is to generate a key string between two distant parties (say, Alice and Bob) such that it remains secret from an eavesdropper (say, Charlie). In the first QKD protocol (BB84) proposed by Bennett and Brassard [31], security is based on the uncertainty of the outcome of incompatible spin measurements chosen randomly along x - and z -directions. The security of standard QKD protocols are based on certain idealistic assumptions [32] that may be minimized in the so-called device independent QKD (DIQKD) [33] where it is no longer required to fully trust the devices used by Alice and Bob. However, practical and loophole free implementations of DIQKD protocols are difficult since they require demonstration of nonlocality. On the other hand, 1s-DIQKD [12] protocols which are intermediate between standard QKD and DIQKD protocols, rely on demonstration of quantum steering for their security. Continuous variable 1s-DIQKD protocols have attracted attention in recent times since they are reasonably robust to losses and practically more feasible compared to their discrete variable counterparts [35].

Monogamy of non-local correlations certifies the security of QKD [34]. To develop a monogamy relation associated with the upper bound of our steering inequality (12), let us consider that three parties Alice, Bob and Charlie share a tripartite state ρ_{ABC} for which the inequality

$$\frac{1}{2}(\Sigma_{BA} + \Sigma_{BC}) \leq \frac{3}{2} \quad (14)$$

is satisfied, where $\Sigma_{BA} = P(b_{\beta_1}|a_{\alpha_1}) + P(b_{\beta_2}|a_{\alpha_2})$ and $\Sigma_{BC} = P(b_{\beta_1}|c_{\gamma_1}) + P(b_{\beta_2}|c_{\gamma_2})$, and c is Charlie's outcome for the measurement chosen from the set

$\{\gamma_1, \gamma_2\}$. The proof comes from contradiction. Let $\frac{1}{2}(\Sigma_{BA} + \Sigma_{BC}) > \frac{3}{2}$. The above sum can be written as the sum of $[P(b_{\beta_1}|a_{\alpha_1}) + P(b_{\beta_2}|c_{\gamma_2})]$ and $[P(b_{\beta_1}|c_{\gamma_1}) + P(b_{\beta_2}|a_{\alpha_2})]$. As the average of the above two terms is greater than $3/2$, one of the terms, say, the first is greater than $3/2$. Then, it is possible to find a conditional Bob's state for which $[P(b_{\beta_1}) + P(b_{\beta_2})] > 3/2$, which contradicts the FUR given by inequality (11). Similarly, using the lower bound of our steering inequality (12), one can obtain $\frac{1}{2}[\Sigma_{BA} + \Sigma_{BC}] \geq \frac{1}{2}$, which together with the relation (14) gives

$$\frac{1}{2} \leq \frac{1}{2}[\Sigma_{BA} + \Sigma_{BC}] \leq \frac{3}{2}. \quad (15)$$

The monogamy relation (14) is used to bound the secret key rate in a 1s-DIQKD protocol. The lower bound of the secret key rate under individual attack is given by [36] $r \geq \mathcal{I}(B : A) - \mathcal{I}(B : C)$, where $\mathcal{I}$ is the mutual information. Suppose that the upper bound of the steering inequality (12) is violated by an amount δ , i.e., $\frac{1}{2}[P(b_{\beta_1}|a_{\alpha_1}) + P(b_{\beta_2}|a_{\alpha_2})] = \frac{3}{4} + \delta$, where $0 < \delta \leq \frac{1}{4}$. Then, the monogamy relation (14) implies that $\frac{1}{2}[P(b_{\beta_1}|c_{\gamma_1}) + P(b_{\beta_2}|c_{\gamma_2})] \leq 0.75 - \delta$. Hence, the lower bound of the key rate becomes

$$r \geq \log_2 \left[\frac{0.75 + \delta}{0.75 - \delta} \right], \quad (16)$$

where the logarithm of base 2 is taken since the secret key rate is expressed in the units of bits per shared state. For the maximally entangled N00N state for which $\delta = 1/4$, the steering inequality (12) is maximally violated, making the lower bound of the secret key rate unity. One may note here that in comparison the lower bound for discrete variables is $1/2$ [14]. Hence, the use of continuous variable systems in QKD offers more security in principle.

V. CONCLUSIONS

To summarize, in the present work we first derive a fine-grained uncertainty relation (FUR) for continuous variable systems with the help of an operational interpretation of the Wigner function [26]. The FUR provides a manifestation of higher uncertainty in continuous variable systems than in discrete variable systems, since in the former the certainty is confined to a lower range bounded by $[1/4, 3/4]$, compared to $[1/2 - 1/(2\sqrt{2}), 1/2 + 1/(2\sqrt{2})]$ for the latter. The increment of uncertainty in continuous variable systems restricts the amount of information leakage to the eavesdropper, making them more secure in principle, than discrete variable systems. Our steering inequality is hence also stronger than that of discrete variable systems, as in the latter, Bob is convinced of the prepared state being entangled only when the average of the conditional probabilities is larger than $1/2 + 1/(2\sqrt{2})$ [14], whereas, in continuous variable systems it is $3/4$. Further, we show that our steering inequality is capable of detecting maximal steerability by

N00N states for $N \geq 2$, thereby circumventing a drawback of the entropic steering inequality which is not violated by the N00N state for $N \geq 2$ [20].

With the help of a derived monogamy relation corresponding to our steering inequality, we bound the key rate in the 1s-DIQKD protocol secured under individual attacks. The relation of the Wigner function with displaced parity operators [26] facilitates comparison of the key rates in continuous and discrete variables. The lower bound of the secret key rate is unity for the shared maximally entangled state of continuous variables, which is double that for discrete variables [14] even when Alice knows Bob's set of observables before preparation of the state. Knowing Bob's set of observables does not help Alice to cheat here, whereas it is indeed helpful in discrete

variable systems [14]. The quasi-probability distributions contained in our steering inequality may be reconstructed experimentally by homodyne detection techniques that are currently realizable with high efficiency [37]. Recent experiments [38] have indeed confirmed Bell violation in continuous variable systems using similar techniques. It should thus be feasible to experimentally verify our steering inequality. Further analysis of more general security attacks, as well as consideration of decoherence effects would be needed to assess the practical viability of such key generation protocols.

Acknowledgements: A.S.M. acknowledges support from the project SR/S2/LOP-08/2013 of DST, India. T.P acknowledges financial support from ANR retour des post-doctorants NLQCC (ANR-12-PDOC-0022- 01).

-
- [1] W. Heisenberg, Z. Phys. **43**, 172 (1927).
 - [2] H. P. Robertson, Phys. Rev. **34**, 163 (1929); E. Schrodinger, Sitzungsberichte der Preussischen Akademie der Wissenschaften, Physikalisch-mathematische Klasse **14**, 296 (1930)..
 - [3] I. Bialynicki-Birula and J. Mycielski, Commun. Math. Phys. **44**, 129 (1975).
 - [4] H. Maassen and J. B. M. Uffink, Phys. Rev. Lett. **60**, 1103 (1988).
 - [5] S. Wehner, A. Winter, New J. Phys. **12**, 025009 (2010).
 - [6] J. Oppenheim and S. Wehner, Science **330**, 1072 (2010).
 - [7] R. Simon, Phys. Rev. Lett. **84**, 2726 (2000); A. Biswas and G. S. Agarwal, New. J. Phys. **7**, 211 (2005); H. Nha, Phys. Rev. A **76**, 014305 (2007); J. Gillet, T. Bastin and G. S. Agarwal, Phys. Rev. A **78**, 052317 (2008).
 - [8] S. Mal, T. Pramanik, A. S. Majumdar, Phys. Rev. A **87**, 012105 (2013).
 - [9] T. Pramanik and A. S. Majumdar, Phys. Rev. A **85**, 024103 (2012); A. Dey, T. Pramanik, and A. S. Majumdar, Phys. Rev. A **87**, 012120 (2013).
 - [10] I. Devetak, & A. Winter, Proc. R. Soc. A **461**, 207 (2005); J. M. Renes, & J. C. Boileau, Phys. Rev. Lett. **103**, 020402 (2009).
 - [11] M. Berta, M. Christandl, R. Colbeck, J. M. Renes, and R. Renner, Nature Phys. **6**, 659 (2010).
 - [12] M. Tomamichel and R. Renner, Phys. Rev. Lett. **106**, 110506 (2011); C. Branciard, E. G. Cavalcanti, S. P. Walborn, V. Scarani, H. M. Wiseman, Phys. Rev. A **85**, 010301(R) (2012).
 - [13] T. Pramanik, P. Chowdhury, A. S. Majumdar, Phys. Rev. Lett. **110**, 020402 (2013).
 - [14] T. Pramanik, M. Kaplan, and A. S. Majumdar, Phys. Rev. A **90**, 050305(R) (2014).
 - [15] F. Grosshans, and P. Grangier, Phys. Rev. Lett. **88**, 057902 (2002); F. Furrer, T. Franz, M. Berta, A. Leverrier, V. B. Scholz, M. Tomamichel, and R. F. Werner, Phys. Rev. Lett. **109**, 100502 (2012).
 - [16] E. Schrödinger, Proc. Cambridge Philos. Soc. **31**, 553 (1935); **32**, 446 (1936).
 - [17] A. Einstein, D. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [18] M. D. Reid, Phys. Rev. A **40**, 913 (1989).
 - [19] S. P. Walborn, A. Salles, R. M. Gomes, F. Toscano, and P. H. Souto Ribeiro, Phys. Rev. Lett. **106**, 130402 (2011).
 - [20] P. Chowdhury, T. Pramanik, A. S. Majumdar, and G. S. Agarwal, Phys. Rev. A **89**, 012104 (2014).
 - [21] Z. Y. Ou, S. F. Pereira, H. J. Kimble, and K. C. Peng, Phys. Rev. Lett. **68**, 3663 (1992).
 - [22] D. J. Saunders, S. J. Jones, H. M. Wiseman, and G. J. Pryde, Nat. Phys. **6**, 845 (2010).
 - [23] B. Wittmann et al., New J. Phys. **14**, 053030 (2012); D. H. Smith et al., Nat. Comm. **3**, 625 (2012); A. J. Bennet, D. A. Evans, D. J. Saunders, C. Branciard, E. G. Cavalcanti, H. M. Wiseman, and G. J. Pryde, Phys. Rev. X **2**, 031003 (2012); V. Handchen et al., Nature Photonics **6**, 598 (2012); S. Steinlechner, J. Bauchrowitz, T. Eberle, and R. Schnabel, Phys. Rev. A **87**, 022104 (2013).
 - [24] J. S. Bell, Physics **1**, 195 (1964); J. F. Clauser, M. A. Horne, A. Shimony, R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
 - [25] H. M. Wiseman, S. J. Jones, and A. C. Doherty, Phys. Rev. Lett. **98**, 140402 (2007).
 - [26] K. Banaszek, and K. Wódkiewicz, Phys. Rev. A **58**, 4345 (1998); Phys. Rev. Lett. **82**, 2009 (1999).
 - [27] H. Lee et al., J. Mod. Opt. **49**, 2325 (2002).
 - [28] G. S. Agarwal, Quantum optics (Cambridge University Press, 2013), Chapter 4.
 - [29] I. Afek, O. Ambar, and Y. Silberberg, Science **328**, 879 (2010).
 - [30] C. F. Wildfeuer, A. P. Lund, J. P. Dowling, Phys. Rev. A **76**, 052101 (2007).
 - [31] C. H. Bennett and G. Brassard, in Proceedings of the IEEE International Conference on Computers, Systems, and Signal Processing, 1984 (unpublished), pp. 31533163.
 - [32] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Rev. Mod. Phys. **74**, 145 (2002); V. Scarani, H. Bechmann-Pasquinucci, N. J. Cerf, M. Dusek, N. Lutkenhaus, and M. Peev, Rev. Mod. Phys. **81**, 1301 (2009).
 - [33] D. Mayers and A. Yao, Proc. of the 39th IEEE Conf. on Foundations of Computer Science (IEEE, Los Alamitos, 1998), p. 503; J. Barrett, L. Hardy, A. Kent, Phys. Rev. Lett. **95**, 010503 (2005); A. Acin et al., Phys. Rev. Lett. **98**, 230501 (2007); N. Gisin, S. Pironio, N. Sangouard, Phys. Rev. Lett. **105**, 070501 (2010). L. Masanes, S. Pironio, and A. Acin, Nat. Comm. **2**, 238 (2011).
 - [34] J. Barrett, N. Linden, S. Massar, S. Pironio, S. Popescu,

- and D. Roberts, Phys. Rev. A **71**, 022101 (2005); J Barrett, A. Kent, and S. Pironio, Phys. Rev. Lett. **97**, 170409 (2006); L. Masanes, A. Acin, N. Gisin, Phys. Rev. A. **73**, 012112 (2006).
- [35] N. Walk, H. M. Wiseman, T. C. Ralph, arXiv: 1405.6593 [quant-ph].
- [36] I. Csiszàr and J. Körner, IEEE Trans. Inf. Theory **24**, 339 (1978).
- [37] K. Wakui, H. Takahashi, A. Furusawa, and M. Sasaki, Optics Express **15**, 3568 (2007).
- [38] A. Kuzmich, I. A. Walmsley, and L. Mandel, Phys. Rev. Lett. **85**, 1349 (2000); H. Jeong, W. Son, M. S. Kim, D. Ahn, and C. Brukner, Phys. Rev. A **67**, 012106, (2003); M. D'Angelo, A. Zavatta, V. Parigi, and M. Bellini, Phys. Rev. A **74**, 052114, (2006).